

# BackRow<sup>®</sup> Max

## DIRECTIONS FOR USE

**IMPORTANT:** This information is approved as part of the Product Label. All instructions within this section must be carefully read in order to obtain safe and successful use of this product.

BackRow Max is an adjuvant designed to reduce the variability in performance of plant protection products applied pre-emergence of the crop.

BackRow Max is authorised for use with all approved herbicides applied in winter and spring sown cereals at full plant protection product rate up to and including G.S. 52 (1/2 inflorescence), and at half the approved rate of the plant protection product thereafter.

### Crop Specific Information

This adjuvant may be used at a maximum concentration of 3.5 % spray solution on the following crops

| Crops / Situation                            | Plant Protection Product  | Max Conc. (% spray sol) | Max No. of Treatments                         | Latest Time of Applic                         |
|--|---|-------------------------|---|---|
| All edible crop production (inc stubble)     | All authorised PPPs at half or less than half their authorised rate | 3.5                     | Follow statutory conditions of use of the PPP | Follow statutory conditions of use of the PPP |
| All non edible crop production (inc stubble) | All authorised PPPs   | 3.5                     | Follow statutory conditions of use of the PPP | Follow statutory conditions of use of the PPP |
| Grass (seed crop)                            | All PPPs authorised for All PPPs authorised for seed production     | 3.5                     | Follow statutory conditions of use of the PPP | Follow statutory conditions of use of the PPP |

In addition, this adjuvant may be used at a maximum concentration of 3.5 % spray solution on a number crops including the following ones, up to the growth stages specified, with all authorised plant protection products applied up to their full authorised rate:

| Crop                                     | Latest Growth Stage Permitted  |
|--|--|
| Table Grapes, Wine Grapes                | Up to and including first fruit set                                      |
| Strawberry                               | Up to and including first fruit development                              |
| Potato                                   | Up to and including tuber initiation                                     |
| Carrot, Parsnip, Turnip, Swede           | Up to and including 9 true leaves unfolded                               |
| Red beet                                 | Up to and including six fully expanded leaves (not for salad-leaf types) |
| Garlic, Bulb Onion, Shallot, Salad Onion | Up to and including 9 leaves clearly visible                             |
| Broccoli/Calabrese                       | Heads begin to form; growing tip > 1 cm                                  |
| Cauliflower                              | Heads begin to form; growing tip up to 1 cm                              |
| Brussels sprout                          | Lateral buds begin to develop  |



| Crop  | Latest Growth Stage Permitted                                     |
|---|---|
| Cabbage   | Heads begin to form   |
| Collard, Kale   | Main shoot 30% typical height                                     |
| Lettuce (nonheading)  | Leaf rosette at 30% typical diameter                              |
| Lettuce (heading)   | Up to heads begin to form   |
| Spinach   | Leaf rosette at 30% typical diameter                              |
| Broad bean (fresh), Dwarf French bean, Runner bean, Edible podded pea | Up to and including first pod set                                 |
| Field bean  | Up to and including first pod set                                 |
| Vining pea  | Up to and including flat pod (non-edible pods only)               |
| Combining pea   | Up to and including flat pod (non-edible pods only)               |
| Soya bean (dry)   | End of flowering; first pods visible (approx. 5 mm) – BBCH 69–609 |
| Linseed   | Up to and including 10% capsules formed                           |
| Oilseed rape  | Up to and including 10% potential pods                            |
| Cereals   | Up to and including GS52 (¼ inflorescence)                        |
| Hops  | Up to cone set  |
| Forage maize  | Up to tip of tassel visible                                       |
| Grain maize   | Up to tip of tassel visible                                       |
| Asparagus   | During dormant phase before BBCH 09                               |

### Restrictions

Please consult Nichino Europe Co., Ltd. or its agents for further details, a full list of approved crops and for specific recommendations. All restrictions of use on the pesticide label must be followed.

### Rates of use

BackRow Max should be used at the following rates, irrespective of water volume:

| Season | Required quantity of BackRow Max |
|--------|----------------------------------|
| AUTUMN | 200 ml/ha                        |
| SPRING | 400 ml/ha                        |

### Mixing

Half fill the spray tank with clean water and commence agitation. Add the recommended quantity of BackRow Max. Then add the recommended quantity of pesticide and the remaining balance of water whilst maintaining agitation throughout.

BACXUK5L25D

BackRow is a registered trade mark of Nichino Europe Co., Ltd

ADJ No. A0978

