

IMPORTANT INFORMATION

FOR USE ONLY AS AN AGRICULTURAL, HORTICULTURAL, INDUSTRIAL AND FORESTRY SPREADING AND WETTING AGENT.

Maximum rate of use: 1.0% of the final spray solution (e.g. 2 litres of product in 200 litres of water per hectare).

READ THE LABEL BEFORE USE. USING THIS PRODUCT IN A MANNER THAT IS INCONSISTENT WITH THE LABEL MAY BE AN OFFENCE. FOLLOW THE CODE OF PRACTICE FOR USING PLANT PROTECTION PRODUCTS.

DIRECTIONS FOR USE

IMPORTANT: This information is approved as part of the Product Label. All instructions within this section must be carefully read in order to obtain safe and successful use of this product.

Sorrento is a unique wetting, deposition and penetration aid designed to improve the retention and uptake of plant protection products and to reduce variability of performance under adverse conditions on a wide range of crops.

Sorrento may be used with **all approved plant protection products** applied at their **full rate** up to and including the following crop growth stages:

Crop	To be applied at or before the following crop growth stages:
Oilseed rape	up to and including 10% potential pods (code 5.1, Sylvester-Bradley, 1985)
Sugar beet	up to and including 6 fully expanded leaves (EPPO code 23)
Cereals	up to and including GS52(1/4 inflorescence)
Potato	up to and including tuber initiation
Forage Maize	up to and including tip of tassel visible (EPPO code 53)
Combining and Vining Peas	up to and including flat pod (EPPO code 205) (non-edible pods only)
Edible podded peas	up to and including pod set (EPPO code 204)

- for use with approved herbicides post-harvest, on stubble and non-crop areas, and for pasture renovation;
- for use with all approved plant protection products applied at half the approved rate in all crops and non-crop situations.

Sorrento can also be used at full rate up to specific growth-stages in a number of other crops, please speak to Nichino Europe Co., Ltd or its agents for specific **recommendations**.

Sorrento is a registered trade mark of Nichino Europe Co., Ltd.

Abate is a registered trade mark of Nichino Europe Co., Ltd.

RATES OF USE

Sorrento should be used at a rate to give a 0.1% dilution of the final spray volume.
For example:

Water Volume	Required Quantity of Sorrento at 0.1%
100 Litres	100ml
200 Litres	200ml

MIXING

Fill the spray tank three quarters full with clean water and commence agitation. Add the recommended quantity of Sorrento. Then add the recommended quantity of pesticide and the remaining balance of water to the spray tank whilst maintaining agitation throughout.

If foaming occurs, add the required quantity of *Abate* to remove foam.

Conditions of supply

All goods supplied by us are of high grade and we believe them to be suitable but as we cannot exercise control over their storage, handling, mixing or use, or the weather conditions before, during and after application, which may affect the performance of the goods, all conditions and warranties, statutory or otherwise, as to the quality of fitness for any purpose of our goods are excluded, and no responsibility will be accepted by us or re-sellers for any failure in performance damage or injury whatsoever arising from their storage, handling, application or use. These conditions cannot be varied by our staff or agents whether or not they supervise or assist in the use of such goods.

